



Maternal Knowledge, Attitudes and Practices towards Prevention and Management of Child Diarrhoea in Urban and Rural Maseru, Lesotho

A. I. Adeleke^{1*} and T. Mhlaba¹

¹*Discipline of Public Health Medicine, School of Nursing and Public Health, University of KwaZulu-Natal, South Africa.*

Authors' contributions

This work was carried out in collaboration among both authors. Author AA designed the study, performed the statistical analysis and wrote the protocol and the first draft of the manuscript. Author MT managed the supervision for conception and design of the study, interpretation of data and drafting of the manuscript. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJTDH/2019/v36i230140

Editor(s):

(1) Dr. Muhammad Chutiyaami, Lecturer, Shehu Sule College of Nursing and Midwifery Damaturu, Nigeria.

Reviewers:

(1) Mahama Bashiru, College of Nursing and Midwifery Nalerigu-North East Region, Ghana.

(2) K. Ramesh Kumar, Kaloji Narayana Rao University of Health Sciences of Health Sciences, India.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/49177>

Received 05 March 2019

Accepted 18 May 2019

Published 27 May 2019

Original Research Article

ABSTRACT

Aim: To compare the knowledge, attitudes and practices of mothers in the prevention and management of child diarrhoea, in rural and urban settings of Maseru.

Methodology: This cross-sectional study was carried out in the Domiciliary (Urban) and Tlali (Rural) Health centres in Maseru, within the period of February to May, 2017. Data was collected from 458 mothers/caregivers, with 299 (65%) and 159 (35%) from urban and rural settings respectively. Scores were assigned for the level of knowledge, attitudes, and practices. STATA 14.1 was applied to determine the strengths of associations between categories of the maternal characteristics and the outcome variables.

Results: Aggregation of participants' knowledge, attitudes, and practices response reveal a statistical significant association with residence. The maternal age range of 30-39 years, $P = .03$, and mothers with three (3) children, $P = .02$ were significantly associated with the knowledge of prevention and management of diarrhoea in the rural area. In the urban area, mothers with tertiary

*Corresponding author: Email: docleke@gmail.com;

education, $P = .04$, employed, $P = .001$, unemployed, $P = .004$, and all categories of monthly income were significantly associated with the knowledge of prevention and management of diarrhoea. For the urban setting, an association between mothers' attitudes and monthly income between M500 – M1399, $P = .05$ was observed. The practices of mothers/caregivers in the prevention and management of diarrhoea showed no significant differences in the light of the socio-demographic variables in both settings.

Conclusion: The study revealed low level of maternal knowledge, attitudes and practices in prevention and management of child diarrhea in the rural and urban settings, hence the need to strengthen the existing health education messages on both settings.

Keywords: Attitude; diarrhoea; knowledge; management; mother; practice; prevention; rural; urban.

DEFINITION FOR THE TERM

- Maternal : Mother of the under-five child, and can be biology or adopted.
Caregiver : Woman that assumes the responsibility of a child in the absence of the biology or adopted mother, which includes; family member or nanny.
Knowledge: The awareness and understanding on the prevention and management of diarrhoea of the under-five child.
Attitude : The condition of readiness for the prevention and management of diarrhoea of under-five child
Practices : The action of performing the process involved in the prevention and management of diarrhoea of under-five child.
Diarrhoea : The passage of three or more loose or watery stools per day, or of more frequency than normal for the individual.
Child : Children aged five years and below.

ABBREVIATIONS

- AIDS : *Acquired Immune Deficiency Syndrome*
CCCD : *Combating Childhood Communicable Diseases*
DALYs : *Disability Adjusted Life Years*
KAP : *Knowledge, Attitude and Practice*
LDHS : *Lesotho Demographic and Health Survey*
MoH : *Ministry of Health*
ORS : *Oral Rehydration Salt Solution*
ORT : *Oral Rehydration Therapy*
SSS : *Sugar and Salt Solution*
UNICEF : *United Nations Children's Fund*
WHO : *World Health Organisation*
SPSS : *Statistical Package for the Social Package*

significant fluid loss, dehydration, and eventually death if the lost fluid is not replaced [4]. The associated risk factors includes; poor hygiene practices, unsafe human waste disposal, lack of safe sanitation, and consuming contaminated drinking water and food [5,6].

Globally in 2010, it was estimated that there were 1.731 billion episodes of diarrhoea in children below the age of five, and about 700 000 of these episodes led to death [7]. The report further indicates that a high proportion of deaths of about 72 per cent occurs in the first 2 years of life [7]. At a prevalence of 22 per cent, about nine (9) per cent of the under-five mortality rate in Lesotho is attributable to diarrhoea infection (UNICEF, 2013). Furthermore, the survey revealed that urban children are slightly less likely to have diarrhoea than rural children (10% versus 13%) [8].

1. INTRODUCTION

Diarrhoeal disease is a major public health problem, that results in illness and death among infants and young children in low-income countries [1]. The disease is caused by the faeco-oral passage of a pathogen through contaminated food or water from the stool of one infected person to the mouth of a new host [2, 3]. This, without prompt attention, may result in

The study in India revealed a tremendous positive improvement in the maternal levels of KAP after the intervention of a structured educational programme on the prevention and management of diarrhoea [9]. The study in Banjul, Gambia, revealed that a mother's high level of knowledge on childhood diarrhoea was associated with parity (number of children), for

mothers of more than two children had a greater knowledge of diarrhoea than mothers of one child. Likewise, mothers with a primary education and above had a better knowledge of diarrhoea when compared to mothers with no formal education [10]. A cross-sectional survey carried out among the rural residents of Sindh in Pakistan revealed poor knowledge on diarrhoea prevention [11]. Only 41 per cent of the respondents identified hand washing as the most important method for prevention of diarrhoea. Furthermore, on prevention, the risk of diarrhoea-related mortality among infants zero to five months of age was higher among those who were partially breastfed at relative risk of 4.62, or not breastfed at relative risk of 10.52 when compared to infants who were breastfed exclusively [12].

The maternal/caregivers' attitudes are important in the fight for the prevention of diarrhoea, as they motivate an individual to adopt a safe healthy practice. The study in Korogocho and Bondo communities of Kenya [13] revealed a positive attitude toward hand washing as most of the urban slum and rural mothers show good attitude towards their hand washing habits on the prevention of infectious diseases, but this was greatly influenced by the availability of water which was only accessible by day.

None persistent diarrhoea could be managed successfully at home with the practise of continuous feeding of the child with diarrhoea, offering more fluids and in the correct and appropriate administration of Oral Rehydration Solution (ORS) [14]. The increase in the child's fluid intake through the use of oral rehydration therapy is the basic intervention for dehydration caused by diarrhoea. Factors such as; the level of knowledge of ORT/ORS, the age of the mothers, availability of prep-packed ORS and education have been found to significantly influence the use of ORT/ORS in the home management of diarrhoea amongst mothers of under-fives [15].

The general notion that the knowledge, attitude and practice (KAP) of mothers in the urban areas is superior to those in the rural areas could be from gathered perceptions and observations, but with no scientific validation. Identifying the gaps in knowledge, attitudes and practices in relation to settings will help plan specific programmes and implement interventions that will reduce the morbidity and mortality associated with diarrhoea.

2. METHODOLOGY

2.1 Study Setting

The research was carried out in two primary healthcare centres, the Domiciliary Clinic in the urban setting and Tlali Health Centre in the rural setting, in Maseru district, Lesotho. Maseru is one of ten districts, with a population of 389,627 people, representing 20.6 per cent of the country's population [16].

2.1.1 Study design and the study population

An observational cross-sectional study was conducted on mothers/caregivers who attended either of the two primary healthcare centres.

2.1.2 Method of selecting sample

Mothers who attended either of the two primary healthcare centres were consecutively enrolled in the study provided they fell within the inclusion criteria at the time of data collection.

2.2 Sample Size

The sample size was determined on the basis of the national under-five diarrhoeal disease prevalence, which is at 22 per cent for Lesotho [8].

The formula for the minimum sample size needed for an interval estimate of a population proportion at 95 per cent confidence interval and five per cent margin of error was:

$$n = \frac{(z\alpha/2)^2 * p * (1-p)}{d^2} = 404$$

Where n is the sample size without the source

$$nf = \frac{n}{1 + \frac{n}{N}} = 404$$

Ten per cent for no response = 40

Total sample size = 444

Where *nf* = final sample size with source

N is the total number of the study population [14,000 (Domiciliary) + 7,221(Tlali clinic) = 21,221].

Based on the proportion of the total population of the two clinics, the Domiciliary and Tlali

health centres will have the sample size of 299 and 159 respectively, totalling 458. In the field, the total number of respondents was 292 and 152 in the urban and rural healthcare centres respectively, making a total of 444 respondents.

2.3 Measurement Instrument and Data Collection Technique

An already existing standardised questionnaire was adapted [10], translated from English to Sesotho (local language), and administered to the participants in the two clinics. Two trained skilled nurses, one in each of the clinic assisted in the filling of the questionnaire.

2.3.1 Data quality assurance

This was achieved through the test and retesting (testing twice) of the questionnaire during the pilot study, on a small scale of ten per cent of the sample size in the two centres.

2.3.2 Data handling/processing

The electronic data was entered into a spreadsheet and coded. The score was assigned for the level of knowledge, attitude, and practice, while STATA 14.1 was used for the regression data analysis. Multiple logistic regression analysis was used to determine the relationship between the dependent variables and socio-demographic variables in the urban and rural settings separately.

2.4 Operational Definitions

Good knowledge: Those mothers who scored three (3) and above from the maximum attainable score of five (5) for the knowledge questions.

Poor knowledge: Those mothers who scored two (2) and below from the maximum attainable score of five (5) for the knowledge questions.

Good attitude: Those mothers who scored three (3) and above from the maximum attainable score of five (5) for the attitude questions.

Poor attitude: Those mothers who scored two (2) and below from the maximum attainable score of five (5) of the attitude questions.

Good practice: Those mothers who scored three (3) and above from the maximum attainable score of five (5) of the practice questions.

Poor Practice: Those mothers who scored two (2) and below from the maximum attainable score of five (5) of the practice questions.

3. RESULTS AND DISCUSSION

3.1 Results

3.1.1 Participants' distribution and profile

A total of 458 mothers/caregivers participated in the study, with urban and rural residents constituting 65 per cent and 35 per cent respectively. The mean age (SD) of respondents was 28.7 (5.1) years in the urban setting, while that of rural setting was 28.4 (9.3) years. In both settings, majority were within the age range of 20 – 29 years (50.8% in urban and 53.5% in the rural). More than half of the respondents were married (60.9% in the urban and 83.6% in the rural setting). Ninety-eight (98.1%) and 24.4 per cents of the respondents in the rural and urban settings respectively were unemployed (Table 1).

3.1.2 Mothers' knowledge on diarrhoeal disease prevention and management

On the mothers'/caregivers' perceived causes of diarrhoea, the majority (93% in urban and 93.7% in rural) of the respondents perceived that unclean water was the main cause of diarrhoea (Fig. 1). The rural respondents did not perceive that specific germs were causes of diarrhoea. Only 6.9 per cent of them mentioned germs as a cause of diarrhoea and 98.1 per cent identified excessive heat (fever) and teething during childhood as causes of diarrhoea.

3.1.3 Mothers' knowledge on the signs of diarrhoea

In the total population of respondents, 10.7 per cent of the respondents in the urban setting, and 32.1 per cent of the rural respondents cited sunken fontanel as a severe form of diarrhoea. Also, only 13 per cent and 31.4 per cent of respondents in the urban and rural respectively recognised that crying without tears is a severe form of diarrhoea. Ninety-nine per cent and 95.6 per cent in the urban and rural settings respectively, were aware that the infection was preventable (Table 2).

Table 1. Percentage distribution of mothers/caregivers place of residence by socio-demographic variables

Socio-demographic variables	Category	Urban (%)	Rural (%)
Mother's Age Group	≤ 19 years	5.4	10.1
	20-29	50.8	53.5
	30-39	43.8	27.0
	≥40	0.0	9.4
	Total	100.0	100.0
Marital status of the mother	Married	60.9	83.6
	Not married	25.4	16.4
	Divorced	13.7	0.0
	Total	100.0	100.0
How many children do you have?	One	35.8	50.9
	Two	49.8	29.6
	Three	12.7	8.2
	Four and above	1.7	11.3
	Total	100.0	100.0
Age of child	0-11	6.7	26.4
	12-35	40.5	55.3
	36-59	52.8	18.2
	Total	100.0	100.0
What is your highest level of education?	Primary	27.4	45.3
	Secondary	60.5	38.4
	Tertiary	9.0	16.4
	Non formal	3.0	0.0
	Other(specify)	0.0	0.0
	Total	100.0	100.0
What is the employment status of mother?	Self-employed	6.8	0.6
	Employed	48.8	1.3
	Unemployed	24.4	98.1
	Total	100.0	100.0
What is the monthly income of the mother?	<500	8.0	0.6
	500-1399	19.4	1.3
	1400-5000	8.7	0.0
	more than 5000	1.7	0.0
	I don't want to say	37.8	0.0
	No income	24.4	98.1
	Total	100.0	100.0

3.1.4 Mothers' attitudes on diarrhoeal disease prevention and management

Maternal attitudes in this context includes their perception on the benefit of exclusive breastfeeding, increased frequency of breastfeeding and administration of fluid during diarrhoea. On the aspect, 100 per cent and 98.3 per cent of the respondents in the rural and urban settings cited the benefit of exclusive breastfeeding in the prevention and management of diarrhoea. Ninety-three per cent and 97.5 per cent of the respondents in the urban and rural settings respectively, agreed that breastfeeding should be increased when the child is infected with diarrhoea (Table 2). In the same vein, 89 per

cent and 98 per cent of the respondents in the urban and rural settings respectively are of good attitudes towards fluid increase for child during diarrhoeal episodes.

3.1.5 Mothers' practices on diarrhoeal disease prevention and management

In terms of the method of prevention adopted by respondents (Table 2), a large percentage (93.6% in the urban and 96.2% in the rural setting) agreed that prevention of diarrhoea in under-five children can be achieved by the washing of hands with soap and water after contact with a child's faecal matter. Only 11.9 per cent of the rural respondents practiced safe and

hygienic preparation of food for diarrhoeal prevention (Table 2). In the urban setting, 64.5 per cent of the respondents gave ORS/SSS as the first line of management to a child with diarrhoea, compared to 8.2 per cent of mothers in the rural setting. The majority of mothers (91.8%) in the rural setting preferred to go to the health centres. On the use of homemade ORS/SSS, in the management of child diarrhoea, this was only practiced by 67.6 per cent and 45.9 per cent of mothers in the urban and rural settings respectively.

3.1.6 Maternal aggregated score of KAP response by residence

Table 3 below revealed that maternal knowledge, attitudes and practices were all statistically significant with residence, at a $P =$ value of .001, .000 and .000 respectively. In all, 77.9, 83.9, and 67.9 per cents of the respondents in the urban setting had good knowledge, attitude and practices respectively as compared to 63.9, 96.9 and 49.1 per cents that had good knowledge, attitude and practices respectively in the rural.




Fig. 1. Maternal knowledge on the causes of diarrhoea

Table 2. Maternal knowledge of symptoms of child diarrhoea, attitudes towards child diarrhoea, prevention and management of child diarrhoea

Variables	Urban (%)	Rural (%)
Knowledge		
Maternal knowledge: Sunken fontanel as a severe form of diarrhoea	10.7	32.1
Maternal knowledge: Cry without tears as a severe form of diarrhoea	13.0	31.4
Maternal knowledge: Diarrhoea is preventable	99.3	95.6
Attitude		
Maternal attitude that exclusive breastfeeding is beneficial	98.3	100
Maternal attitude to breastfeed increase for child with diarrhoea	93	97.5
Maternal attitude to fluid increase for child with diarrhoea	89	98
Preventative and management practices		
Mother's preventative practices for child diarrhoea		
Washing your hands with soap and water after being in contact with a child's faecal matter	93.6	96.2
Safe and hygienic preparation of food	96.0	11.9
Safe and hygienic disposal of faecal and contaminated materials	20.7	9.4
Mother's first line of management for child diarrhoea		
Give oral rehydration solution or ready-made sachets	64.5	8.2
Go to a health centre	23.4	91.8
Maternal use of homemade ORS/SSS	67.6	45.9
Maternal correct mixing of homemade ORS/SSS	64.9	5.7

Table 3. Maternal/caregiver's knowledge, attitude, and practices (KAP) by residence

Maternal KAP		Residence		Pearson Chi-square test
		Rural (%)	Urban (%)	
Knowledge	Good knowledge	63.5	77.9	.001
	Poor knowledge	36.5	22.1	
Attitude	Good attitude	96.9	83.9	.000
	Poor attitude	3.1	16.1	
Practice	Good practice	49.1	67.9	.000
	Poor practice	50.9	32.1	

3.1.7 Associated factors of maternal knowledge, attitudes and practice in relation to residence

In the rural setting (Table 4). Mothers/caregivers who were 30-39 years were approximately 79 per cent less likely to have good knowledge as compared to mothers below the age of 19 years (AOR: .21, $P = .03$). Mothers with three (3) children were approximately eight times more likely to have good knowledge as compared to those with one child (AOR: 7.66, $P = .02$).

In the urban setting (Table 5). Mothers/caregivers with a tertiary education were 72 per cent less likely to have good knowledge as compared to mothers with a primary education (AOR: .28, $P = .04$). Employed mothers/caregivers were approximately 80 per cent less likely to have good knowledge as compared to the self-employed mothers (AOR: .17, $P = .001$). Unemployed mothers/caregivers were approximately six times more likely to have good knowledge as compared to the self-employed mothers (AOR: 5.56, $P = .004$). Mothers/caregivers with a monthly income of M500-M1319 (AOR: 5.97, $P = .01$), M1400-M5000 (AOR: 9.37, $P = .003$), and more than M5000 (AOR: 23.35, $P = .01$) were six, nine and 23 times respectively, more likely to have good knowledge as compared to the mothers/caregivers with a monthly income of less than M500.

Mothers/caregivers with non-formal education were 82% less likely to have good attitude as compared with mothers with primary education (AOR: .18, $P = .05$). Mothers with a monthly income of between M500-M1399 were four times more likely to have a good attitude as compared to mothers with a monthly income of less than M500 (AOR: 3.9, $P = .05$).

3.2 Discussion

3.2.1 Demographic profile

In this study, mothers/caregivers in the age group of 20 -29 years constituted the majority of the respondents in both settings. A study in India [9] and Pakistan [17], reported similar finding, for the highest number of respondents were within the age group of 20 - 29 years. The rural respondents had higher numbers of children (parity), at four children and above compared to urban respondents. This finding is congruent with the characteristics of households in Lesotho [16].

In the study, highest number of respondents completed secondary school level in the urban setting as compared to the rural settings with highest percentage of respondents completing primary school level. The lower percentage recorded in the rural setting on secondary school level could partly be as a result of the limited educational facilities particular to the rural setting on this cadre. There were more employed respondents in the urban, as compared to the rural setting. Therefore, the majority of respondents in the rural setting had no source of income. This was in tandem with the demographic survey previously conducted in Lesotho [8], with a higher employment status in the urban setting compared to the rural settings.

3.2.2 Respondents' knowledge on prevention and management of diarrhoea (Fig. 1 and Tables 2 and 3)

On both settings, practically all respondents agreed that unclean water contributed to diarrhoea, results which were contrary to the study conducted in Kanyakumari district, South India [18], where only four per cent of respondents identified unclean water as a cause of diarrhoea. The knowledge about the association between diarrhoea being caused by

Table 4. Multiple logistic regression analysis of socio-demographic characteristics and maternal KAPs on diarrheal prevention and management in under-five children (rural residence)

Demographic characteristics		Knowledge				Attitude				Practice			
		Poor (%)	Good (%)	AOR(CI)	P-Value	Poor (%)	Good (%)	AOR(CI)	P-Value	Poor (%)	Good (%)	AOR(CI)	P-Value
Age of the mother or caregiver (years)	< 19	4(25.0%)	12(75%)	--	--	1(6.3%)	15(93.8%)	--	--	9(56.3%)	7(43.8%)	--	--
	20-29	29(34.1%)	56(65.9)	0.58(0.17,2.06)	.40	3(3.5%)	82(96.5%)	2.74(0.16,46.2)	.49	48(56.5%)	37(43.5%)	0.92(0.30,2.85)	.89
	30-39	20(46.5%)	23(53.5%)	0.21(0.05,0.88)	.03	1(2.3%)	42(97.7%)	7.14(0.21,248)	.28	20(46.5%)	23(53.5%)	1.28(0.34,4.77)	.72
	40 +	5(33.3%)	10(66.7%)	0.20(0.03,1.33)	.10	0(0%)	15(100.0%)	1	-	4(26.7%)	11(73.3%)	2.82(0.47,16.9)	.26
Marital status of the mother	Married	48(36.1%)	85(63.9%)	--	--	3(2.3%)	130(97.7%)	--	--	66(49.6%)	67(50.4%)	--	--
	Single (Not married)	10(38.5%)	16(61.5%)	0.88(0.34,2.23)	.78	2(7.7%)	24(92.3%)	0.56(0.07,4.82)	.60	15(57.7%)	11(42.3%)	0.87(0.36,2.12)	.76
	Divorced	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-
Number of children	One	31(38.3%)	50(61.7%)	--	--	2(2.5%)	79(97.5%)	--	--	46(56.8%)	35(43.2%)	--	--
	Two	20(42.6%)	27(57.4%)	1.20(0.51,2.79)	.68	3(6.4%)	44(93.6%)	0.13(0.01,1.31)	.08	23(48.9%)	24(51.1%)	1.19(0.53,2.67)	.68
	Three	2(15.4%)	11(84.6%)	7.66(1.32,44.6)	.02	0(0.0%)	13(100%)	1	-	5(38.5%)	8(61.5%)	1.39(0.36,5.44)	.64
	Four +	5(27.8%)	13(72.2%)	4.05(0.92,17.8)	.06	0(0.0%)	18(100%)	1	-	7(38.9%)	11(61.1%)	1.09(0.28,4.22)	.90
Education level	Primary	26(36.1%)	46(63.9%)	--	--	0(0%)	72(100.0%)	1	--	33(45.8%)	39(54.2%)	--	--
	Secondary	21(34.4%)	40(65.6%)	1.27(0.58,2.77)	.55	3(4.9%)	58(95.1%)	2.55(0.28,23.1)	.41	32(52.5%)	29(47.5%)	0.81(0.39,1.67)	.57
	Tertiary	11(42.3%)	15(57.7%)	0.95(0.35,2.66)	.92	2(7.7%)	24(92.3%)	1	-	16(61.5%)	10(38.5%)	0.65(0.24,1.76)	.40
	Non-formal	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-
	Other(specify)	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-
Employment status	Self Employed	1.0(100%)	0(0%)	1	--	0(0%)	1(100.0%)	1	-	0(0%)	1(100.0%)	1	-
	Employed	1.0(50%)	1(50.0%)	0.73(0.04,25.2)	.83	0(0%)	2(100.0%)	1	-	0(0%)	2(100.0%)	1	-
	Unemployed	56(35.9%)	100(64.1%)	1	-	5(3.2%)	151(96.8%)	1	-	81(51.9%)	75(48.1%)	1	-
	Pensioner	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-
	Receiving Disability Grant	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-
Monthly income (Maloti)	Other(specify)	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-
	< 500	1.0(100%)	0(0%)	1	-	0(0%)	1(100.0%)	1	-	0(0%)	1(100.0%)	1	--
	500-1399	1.0(50%)	1(50.0%)	1	-	0(0%)	2(100.0%)	1	-	0(0%)	2(100.0%)	1	-
	1400-5000	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-
	> 5000	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-
	I don't want to say	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-
No income	56(35.9%)	100(64.1%)	1	-	5(3.2%)	151(96.8%)	1	-	81(51.9%)	75(48.1%)	1	-	

*AOR = Adjusted Odd Ratio, CI = Confidence Interval; -- = reference category; - = omitted/ empty

Table 5. Multiple logistic regression analysis of socio-demographic characteristics and maternal KAPs on diarrheal prevention and management in under-five children (Urban residence)

Demographic characteristics		Knowledge				Attitude				Practice			
		Poor (%)	Good (%)	AOR(CI)	P-Value	Poor (%)	Good (%)	AOR(CI)	P-Value	Poor (%)	Good (%)	AOR(CI)	P-Value
Age of the mother or caregiver (years)	< 19	2(12.5%)	14(87.5%)	--	--	5(31.3%)	11(68.8%)	--	--	6(37.5%)	10(62.5%)	--	--
	20-29	33(21.7%)	119(78.3%)	0.66(0.11,4.08)	.66	23(15.1%)	129(84.9%)	2.84(0.78,10.4)	.12	47(30.9%)	105(69.1%)	1.04(0.33,3.31)	.94
	30-39	31(23.7%)	100(76.3%)	0.56(0.09,3.57)	.54	20(15.3%)	111(84.7%)	2.75(0.70,10.8)	.15	43(32.8%)	88(67.2%)	0.85(0.26,2.78)	.79
	40 +	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-
Marital status of the mother	Married	43(23.6%)	139(76.4%)	--	--	32(17.6%)	150(82.4%)	--	--	54(29.7%)	128(70.3%)	--	--
	Single (Not married)	16(21.1%)	60(78.9%)	1.00(0.45,2.22)	1.00	11(14.5%)	65(85.5%)	1.27(0.54,2.98)	.59	26(34.2%)	50(65.8%)	0.96(0.51,1.81)	.91
	Divorced	7(17.1%)	34(82.9%)	1.43(0.51,4.08)	.50	5(12.2%)	36(87.8%)	1.63(0.54,4.97)	.39	16(39.0%)	25(61.0%)	0.74(0.35,1.58)	.44
Number of children	One	25(23.4%)	82(76.6%)	--	--	19(17.8%)	88(82.2%)	--	--	36(33.6%)	71(66.4%)	--	--
	Two	28(18.8%)	121(81.2%)	1.46(0.69,3.09)	.32	20(13.4%)	129(86.6%)	1.28(0.56,2.91)	.56	51(34.2%)	98(65.8%)	0.92(0.51,1.66)	.78
	Three	11(28.9%)	27(71.1%)	0.70(0.25,2.00)	.51	8(21.1%)	30(78.9%)	0.61(0.20,1.89)	.39	8(21.1%)	30(78.9%)	2.30(0.87,6.08)	.09
	Four +	2(40.0%)	3(60.0%)	0.27(0.04,1.95)	.19	1(20.0%)	4(80.0%)	0.51(0.05,5.43)	.57	1(20.0%)	4(80.0%)	2.32(0.23,23.8)	.48
Education level	Primary	19(23.2%)	63(76.8%)	--	--	15(18.3%)	67(81.7%)	--	--	25(30.5%)	57(69.5%)	--	--
	Secondary	32(17.7%)	149(82.3%)	1.44(0.66,3.12)	.36	21(11.6%)	160(88.4%)	1.41(0.64,3.11)	.40	63(34.8%)	118(65.2%)	0.86(0.47,1.58)	.63
	Tertiary	11(40.7%)	16(59.3%)	0.28(0.09,0.92)	.04	8(29.6%)	19(70.4%)	0.37(0.11,1.27)	.11	7(25.9%)	20(74.1%)	1.17(0.40,3.48)	.77
	Non- formal	4(44.4%)	5(55.6%)	0.74(0.15,3.76)	.72	4(44.4%)	5(55.6%)	0.18(0.03,1.03)	.05	1(11.1%)	8(88.9%)	2.47(0.26,23.5)	.43
	Other(specify)	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-
Employment status	Self Employed	12(15%)	68(85.0%)	--	--	10(12.5%)	70(87.5%)	--	--	22(27.5%)	58(72.5%)	--	--
	Employed	45(30.8%)	101(69.2%)	0.17(0.06,0.46)	.001	24(16.4%)	122(83.6%)	0.69(0.26,1.80)	.45	46(31.5%)	100(68.5%)	0.77(0.39,1.50)	0.44
	Unemployed	9(12.3%)	54(87.7%)	5.56(1.71,18.0)	.004	14(19.2%)	59(80.8%)	1.76(0.55,5.63)	.34	28(38.4%)	45(61.6%)	0.41(0.13,1.27)	.12
	Pensioner	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-
	Receiving Disability Grant	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-
	Other(specify)	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-	0(0%)	0(0%)	-	-
Monthly income	Less than 500	13(54.2%)	11(45.8%)	--	--	7(29.2%)	17(70.8%)	--	--	5(20.8%)	19(79.2%)	--	--
	500-1399	17(29.3%)	41(70.7%)	5.97(1.73,20.6)	.01	8(13.8%)	50(86.2%)	3.90(1.02,14.9)	.05	12(20.7%)	46(79.3%)	1.06(0.31,3.62)	.93
	1400-5000	9(34.6%)	17(65.4%)	9.37(2.14,41.0)	.003	5(19.2%)	21(80.8%)	2.85(0.60,13.7)	.19	5(19.2%)	21(80.8%)	1.34(0.30,5.93)	.70
	more than 5000	2(40.0%)	3(60.0%)	23.35(2.04,267)	.01	2(40.0%)	3(60.0%)	2.33(0.21,25.4)	.49	0(.0%)	5(100.0%)	1	-
	I don't want to say	16(14.2%)	97(85.8%)	21.35(5.94,76.7)	.00	12(10.6%)	101(89.4%)	4.43(1.31,15.0)	.02	46(40.7%)	67(59.3%)	0.45(0.15,1.37)	.16
	No income	9(12.3%)	64(87.7%)	1	-	14(19.2%)	59(80.8%)	1	-	28(38.4%)	45(61.6%)	1	-

*AOR = Adjusted Odd Ratio, CI = Confidence Interval; -- = reference category; - = omitted/ empty

germ was poor in the rural setting, where almost all respondents cited teething as the main cause of diarrhoea. On the contrary, in the urban setting, virtually all respondents cited germs as a cause of diarrhoea, but two third of the respondents likewise cited teething as a possible cause (Fig. 1). This is supported by the study conducted in North of Saudi Arabia, where three quarter of the informants cited teething as a possible cause of diarrhoea [19]. On severity, only 11 and 32 per cents of respondents in the urban and rural settings respectively recognised sunken fontanel as severe sign of diarrhoea. This is similar to the study in the rural community in Kenya where only 3.1 per cent [14] of the respondents recognised sunken fontanel as severe sign of diarrhoea. In terms of prevention, respondents on both settings had good knowledge that diarrhoeal disease was preventable.

In the study, on the average, the level of good knowledge on the prevention and management of diarrhoea was higher among the urban respondents as compared to the rural respondents (Table 3). This can be linked to the observation made in the study conducted in Ghana, where the risk of childhood diarrhoea was found to be significantly higher in rural areas than urban areas [20]. Though the level of knowledge in the prevention and management of diarrhoea in the two settings was above average (higher in the urban). This will be considered insufficient when related to the study carried out in Pakistan [21] where the level of respondents' knowledge was 75 per cent. The difference in the level of knowledge observed in this study between the two settings, might have been due to the access of more information on the prevention and management of diarrhoea in the urban setting, secondary to the availabilities of more healthcare facilities in this setting as compared to the rural setting.

3.2.3 Respondents' attitudes to prevention and management of diarrhoea (Tables 2 and 3)

The study revealed that, the respondents in the rural and urban settings had good attitudes to breastfeeding and recommended that it be increased in the event that a child has diarrhoea. This was in line with a study done in Kosovo where more than 75 per cent of respondents breastfed their babies more than usual during episodes of diarrhoea [22]. Similarly, increased

fluid intake was believed to benefit a child in the case of diarrhoea, which was affirmed by the respondents in both settings. This was however contrary to a previous health survey conducted in Lesotho where 28.8 per cent of urban and 19.3 per cent of rural respondents believed that increased fluid intake was beneficial during diarrhoea [8]. The difference between this study's findings and the Lesotho Health Survey (LDHS) may be due to recent improvements in attitudes due to information received on this aspect. Furthermore, it may be due to differences in population size, settings and methods adopted in the LDHS study. Respondents on both settings had good attitudes towards exclusive breastfeeding, but with higher percentage in the rural setting. These findings were however contrary to findings of a study conducted in Anantapur district in India, where only 8.9 per cent of the respondents had good attitudes to exclusive breastfeeding [9]. This could be due to the fact that higher numbers of rural respondents were unemployed, and could afford more time with their children, therefore increasing the chances to exclusively breastfeed their children as compared to their urban counterparts.

3.2.4 Respondents' practices on prevention and management of diarrhoea (Tables 2 and 3)

There was high level of good practice on both settings on washing of hands with water and soap when hands were contaminated with a child's faecal matter. The importance of this practice was revealed in the conclusions drawn from systemic review of literatures in the low- and middle-income countries in Asia, where diarrheal morbidity was reduced by one-third through hand washing interventions [23]. The study revealed a poor practice in the rural setting on the part of safe and hygienic preparation of food as a useful exercise in the prevention and management of diarrhoea, unlike in the urban. This was in line with findings in rural Soweto, South Africa where only 1.5 per cent of the respondents practiced hygienic practices such as the washing of utensils and bottles when preparing SSS [24]. Reasons could be that there may be assumptions that utensils were generally clean therefore need no further washing with water and soap before use.

In the management of diarrhoea, two-third of the urban respondents cited the use of oral rehydration solution or ready-made sachets as

the first line of management, while less than a tenth of the rural respondents agreed to similar practice. A study conducted in Asia and Africa, reported only a fifth of caretakers who gave their children ORS [25]. Majority of the respondents in the rural settings preferred going to health centres as the first line of management, unlike in the urban, where minority (a fifth) cited similar practice. Only two-third of the urban respondents used homemade oral rehydration solution, while less than half of the rural respondents had similar practice. This is similar to the study in Kanyakumari district, south India, where only 50 per cent of respondents prepared ORS at home [18]. In terms of the correct mixing of SSS, less than a tenth of rural respondents knew how to correctly mix the solution, unlike in the urban, where two-third can correctly do proper mixing of the solution. This was similar to a study done in Johannesburg, South Africa, where only 21 per cent of mothers/caregivers correctly prepared homemade ORS [24]. Other studies in Nigeria [26] and India [27] similarly reported these findings in like manner.

3.2.5 Factors associated with KAP outcome in the prevention and management of diarrhoea (Tables 4 and 5)

In the rural setting, age and the number of children (parity) by the respondents were significant factors (predictors) for their knowledge on the prevention and management of diarrhoea. The older age category of 30-39 years is more at risk on good knowledge as compared to the reference category (≤ 19 years), and this might be due to the fact that the older mothers might not have had access to formal education, unlike the younger ones. Likewise, mothers with three (3) children were more likely to have good knowledge as compared to those with one child. It can be inferred that such mothers with a high parity had more experience and prior knowledge and understanding of diarrhoea prevention and management. This is similar to studies carried out in Gambia [10] and Nepal [28].

In the urban setting, educational level, employment status, and monthly income were predictors in the prevention and management of diarrhoea in under-five children. Mothers/caregivers with tertiary education were less likely to have good knowledge as compared to those with a primary education. This is contrary to the study in Saudi Arabia, where, it was identified that the knowledge of mothers improves with education [19]. The difference

might have been due to the differences in the study population and setting. Also, mothers/caregivers who were unemployed were more likely to have good knowledge in relation to self-employed mothers. This may be due to the unlimited time and undivided attention possessed by this category (unemployed) to seek knowledge. Higher monthly income was associated with a significant increase in the level of knowledge when compared to those who earned less than M500. This was supported by a study in Ghana, where the odds of diarrhoea incidence were significantly higher among the rural poorer respondents [20]. Furthermore, monthly income was found to be a predictor in the level of attitude in the prevention and management of diarrhoea. In the urban setting, mothers with a monthly income of between M500 – M1399 were more likely to have a good attitude as compared with those who earned less than M500.

3.2.6 Challenges and limitations

A cross-sectional nature of this study, subjects it to biases when determining the associations of independent variables to dependent ones. However, measures to reduce bias in this study were employed, such as in the use of a standardised questionnaire and in the training of the research assistants, however, the representativeness of this study sample to the population was not guaranteed. The sample only represented participants in the two centres and not the entire district, therefore basing the study on a larger sample size could have generated more accurate or stronger results.

4. CONCLUSION AND RECOMMENDATIONS

4.1 Conclusion

It was established that, there were differences in the knowledge, attitudes and practices in the two settings. The findings of the study further revealed that various socio-demographic characteristics in both the urban and rural settings influenced, particularly, maternal knowledge on prevention and management of child diarrhoea. Monthly income was solely associated with maternal attitudes in the urban setting. In addition, there were no observed significant influences of socio-demographic characteristics on maternal practices in both settings.

4.2 Recommendations

Based on the findings of the study, it is recommended that: There is the need to strengthen health education messages on childhood diarrhoea. This ought to be in a form of a repeated structured educational programme to educate mothers/caregivers, basically on the causes, signs and severity of diarrhoea in settings such as schools, hospitals/clinics, and other work areas. Though mothers/caregivers in the urban setting were more familiar with methods of prevention of diarrhoea in children under the age of five, but need to be re-iterated in both settings. For better coverage of health messages, these should be disseminated through clinic visitations by health officials, media platforms and community campaigns. The use of ORS and/or homemade SSS as the first line of management of diarrhoea, should be emphasised in both settings, as uncomplicated diarrhoea can be successfully managed at home. In addition, the importance of ORS/homemade SSS as a lifesaving intervention and the correct mixing of the solution should be taught.

Further research using a more rigorous study designs that involve the combination of quantitative and qualitative (in-depth) research methodologies, and a larger sample size may provide stronger evidence in addressing the research question.

CONSENT AND ETHICAL APPROVAL

This was according to the Biomedical Research Ethics Committee (BREC) of UKZN and the Ethics Committee of the Ministry of Health, Lesotho. The participant consent and ethical approval were collected and preserved by the authors.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kotloff KL, Nataro JP, Blackwelder WC, Nasrin D, Farag TH, Panchalingam S, et al. Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multicenter Study, GEMS): A prospective, case-control study. *The Lancet*. 2013; 382(9888):209-22.
2. Walker CLF, Perin J, Aryee MJ, Boschi-Pinto C, Black RE. Diarrhea incidence in low-and middle-income countries in 1990 and 2010: a systematic review. *BMC Public Health*. 2012;12(1):220.
3. Kelly P. Diarrhoeal disease. *Journal of the Royal College of Physicians*. 2011;11(5): 488-91.
[Accessed 12 May 2019]
Available: <http://www.clinmed.rcpjjournal.org/content/11/5/488.full.pdf>
4. Unicef. Diarrhoea: why children are still dying and what can be done.
Available: http://www.unicef.org/media/files/Final_Diarrhoea_Report_October_2009_final.pdf. 2010:68
[Accessed 23 May 2018]
Available: https://apps.who.int/iris/bitstream/handle/10665/44174/9789241598415_eng.pdf;sequence=1
5. Kahabuka C, Kvale G, Hinderaker SG. Factors associated with severe disease from malaria, pneumonia and diarrhea among children in rural Tanzania—A hospital-based cross-sectional study. *BMC infectious diseases*. 2012;12(1):219.
6. Peter G, Nkambule SE. Factors affecting sustainability of rural water schemes in Swaziland. *Physics and Chemistry of the Earth, Parts A/B/C*. 2012;50:196-204.
7. Walker CLF, Rudan I, Liu L, Nair H, Theodoratou E, Bhutta ZA, et al. Global burden of childhood pneumonia and diarrhoea. *The Lancet*. 2013;381(9875): 1405-16.
8. Ministry of Health. *Child Health*. In: Ministry of Health, editor. *Lesotho Demographic and Health Survey*. 27-07-2016 ed. Rockville, MD 20850, USA; ICF International. 2014;538.
9. Sunanda G, Ramaiah D, Sadiq MMJ, Narayana G. Impact of structured educational program on maternal knowledge, attitude, and practice toward diarrhea management in children < 5 years age in Anantapur District. *CHRISMED Journal of Health and Research*. 2017; 4(3):186.
10. Sillah F. Knowledge, attitude and practice assessment of mothers in managing children under five years old with diarrhea in the Gambia. *Taipei Medical University Health Nutrition 1st Division*. 2012;1-71.
11. Khan AI, Danish SH, Ashfaq A, Ahmad F, Mohammad S, Warsi A, et al. Knowledge, attitude and practices of mothers regarding diarrheal risk factors and management in

- under 5 children: A cross sectional survey in Dadu and Badin Districts of Sindh, Pakistan. *Journal of Dow University of Health Sciences*. 2016;10(1).
12. Salone LR, Vann Jr WF, Dee DL. Breastfeeding: An overview of oral and general health benefits. *The Journal of the American Dental Association*. 2013;144(2): 143-51.
 13. USAID. Handwashing Knowledge, Attitudes, and Practices in Korogocho (Nairobi) and Bondo (Nyanza) The Maternal and Child Health Integrated Program (MCHIP). 2014;30.
 14. Othero DM, Orago AS, Groenewegen T, Kaseje DO, Otengah P. Home management of diarrhea among underfives in a rural community in Kenya: Household perceptions and practices. *East African Public Health Association*. 2009;5(3):5. [Accessed 23 May 2018] Available:<https://tspace.library.utoronto.ca/bitstream/1807/39197/1/lp08026.pdf>
 15. Bello DA, Afolaranmi TO, Hassan ZI, Ogbonna FC, Inedu PG, Ejiga C, et al. Knowledge and use of oral rehydration solution in the home management of diarrhea among mothers of under fives in Jos, Plateau State. *International Journal of Biomedical Research*. 2017;8(1):5.
 16. Lesotho Bureau of Statistics. Lesotho Demographic Survey 2011. In: Ministry of Development Planning GoL, editor. Lesotho. 2011;367.
 17. Asghar I, Javaid A, Anwar M, Mustafa G. Knowledge and practices of rural mothers about diarrhea among infants. *Journal Of Sheikh Zayed Medical College*. 2016; 7(4):1053-6.
 18. Nelson SB, Panicker PR, Nandakumar L, Malaichamy K. Knowledge and practice regarding diarrheal diseases and drinking water usage in Kanyakumari district, South India. *National Journal of Research in Community Medicine*. 2017;6(3):217-222.
 19. Alshammari BA, Moussa S, Zaid AW. Knowledge and behavioral practice of mothers about diarrhea in Hail region, Saudi Arabia. *World Journal of Pharmaceutical Research*. 2017;7(3):1661-74. [Accessed 04 July 2018] Available:http://www.wjpr.net/dashboard/abstract_id/9072
 20. Kumi-Kyereme A, Amo-Adjei J. Household wealth, residential status and the incidence of diarrhoea among children under-five years in Ghana. *Journal of Epidemiology and Global Health*. 2016;6(3):131-40.
 21. Zafar M. Knowledge and attitude towards and preventive practices relating to diarrhea among mothers under five years of Children: Findings of a cross-sectional study in Karachi. *Pakistan J Infect Dis Ther*. 2014;2(1):1-6.
 22. Berisha M, Hoxha-Gashi S, Gashi M, Ramadani N. Maternal practice on management of acute diarrhea among children under five years old in kosova. *TAF Preventive Medicine Bulletin*. 2009; 8(5).
 23. Shah D, Choudhury P, Gupta P, Mathew JL, Gera T, Gogia S, et al. Promoting appropriate management of diarrhea: A systematic review of literature for advocacy and action: UNICEF-PHFI series on newborn and child health, India. *Indian pediatrics*. 2012;49(8):627-49.
 24. Llale MA. How well do mothers and caregivers prepare oral rehydration solution. *Wits Institutional Repository environment on DSpae: University of the Witwatersrand medical school*; 2017. [Accessed 04 July 2018] Available:<http://wiredspace.wits.ac.za/bitstream/handle/10539/23112/Andrew%20Llale-2016.pdf?sequence=1>
 25. Nasrin D, Wu Y, Blackwelder WC, Farag TH, Saha D, Sow SO, et al. Health care seeking for childhood diarrhea in developing countries: evidence from seven sites in Africa and Asia. *The American Journal of Tropical Medicine and Hygiene*. 2013;89(Suppl_1):3-12.
 26. Adimora G, Ikefuna A, Ilechukwu G. Home management of childhood diarrhoea: Need to intensify campaign. *Nigerian Journal of Clinical Practice*. 2011;14(2):237-41.
 27. Mahor G. Knowledge and attitudes of mothers regarding use of oral rehydration solution in management of diarrhea. *Asian Journal of Biomedical and Pharmaceutical Sciences*. 2013;3(22):6.
 28. Ansari M, Palaian S, Ibrahim M. The role of mothers in the management of childhood diarrhoea in Nepal. *Australasian Medical Journal*. 2009;1(14):235-8.

List of Appendices

1. Data collection tool (questionnaire)
2. Informed consent and participants' declaration
3. Lesotho ministry of health approval
4. Brec (Biomedical research ethics committee) approval

APPENDIX ONE: DATA COLLECTION TOOL

Part 1: Demographic information

1. Age of mother:

No	Questions and filters	Coding categories	Skip to
2	Marital status	Married.....1 Not married.....2 Divorced.....3	
3	How many children do you have?	One.....1 Two.....2 Three.....3 Others please specify.....4	
4	Age of child / age of last child if more than one	
5	What is your highest level of education?	Primary.....1 Secondary.....2 Tertiary.....3 Non-formal.....4 Other, please specify.....5	
6	Employment status	Self-employed.....1 Employed.....2 Unemployed.....3 Pensioner.....4 Receiving disability grant.....4 Other, please specify.....5	
7	Monthly Income?	Less than M500.....1 Between M500-M1399.....2 Between M1400-M5000.....3 More than R5000.....4 I don't want to say.....5	Between

Part 2: Knowledge on diarrhoea

No	Questions and filters	Coding categories	Skip to
8	What do you think are the causes of diarrhoea? (you may choose more than one answer)	Unclean water.....1 Dirty hands.....2 Germs.....3 Over eating.....4 Excessive heat (environmental).....5 Excessive cold (environmental).....6 Teething.....7 Side effects of certain medications.....8 Others, please specify.....9	
9	Does breastfeeding increase the risk of child diarrhoea?	Yes.....1 No.....2 Don't know.....3	

No	Questions and filters	Coding categories	Skip to
10	Can diarrhoea be caused by prolonged breast feeding that lasts up to two years?	Yes.....1 No.....2 Don't know.....3	
11	Infant formula feeding can pose a higher risk of diarrhoea compared to breast feeding?	Yes.....1 No.....2 Don't know.....3	
12	Can diarrhoea be life-threatening?	Yes.....1 No.....2 Don't know.....3	
13	Where/from whom do you/ did you receive information on prevention and management of child diarrhoea?	Never.....1 Relative.....2 Friend.....3 Health worker.....4 Work shop.....5 Others, please specify.....6 Don't know.....7	
14	How do you know if your child has diarrhoea?	Passage of normal stool at least twice a day.....1 Passage of three or more loose or watery stools per day.....2 Others, please specify.....3 Don't know.....4	

Part 3: Attitudes and feeding practices towards diarrhoea and its management

No	Questions and filters	Coding categories	Skip to
15	Do you believe in exclusive breast feeding? (for nursing mothers)	Yes.....1 No.....2 Don't know.....3	
16	How often should breastfed babies be best fed?	On demand.....1 1 - 2 times daily.....2 3 times daily.....3 Don't know.....4	
17	In the presence of child diarrhoea, what should a breastfeeding mother do?	Do not breast feed.....1 Reduce breast feeding.....2 Increase breast feeding.....3 Breastfeed normally as when not with diarrhoea.....4 Others, specify.....5 Don't know.....6	
18	What is the most beneficial duration of breastfeeding?	Less than 6 months.....1 6 – 12 months.....2 Greater than 12 months.....3	
19	What complimentary foods do you give your child when introduced to solids?	Rice and sauce.....1 Pap only.....2 Pap plus other supplementary foods.....3 Other, specify.....4	
20	Do you think that more liquids should be given to a child with diarrhoea?	Yes.....1 No.....2 Don't know.....3	

Part 4: Practice towards diarrhoea disease prevention

No	Questions and filters	Coding categories	Skip to
21	Do you believe that diarrhoea is preventable? (if no, please go to question 26)	Yes.....	1
		No.....	2
		Don't know.....	3
22	If yes, how do you prevent it? (you may choose more than one answer)	Washing your hands with soap and water after getting contact with a child's faecal matter.....	1
		Safe and hygienic preparation of food.....	2
		Safe and hygienic disposal of faecal and contaminated materials.....	3
		Others, please specify.....	4
		Pit latrine.....	1
23	What sanitation facilities do you use at home?	Flush toilet.....	2
		Openly defecate in the compound premises.....	3
		Bed pan/ Potty.....	4
24	What is your source of drinking water?	River water.....	1
		Open well.....	2
		Public tap.....	3
		Private tap.....	4
		Borehole.....	5
		Other.....	6

Part 5: Practices towards diarrhoea disease management

No	Questions and filters	Coding categories	Skip to
25	What would you do when your child has diarrhoea?	Do nothing.....	1
		Go to a health centre.....	2
		Give oral rehydration solution or ready-made sachets.....	3
		Use traditional herbs.....	4
		Go to traditional herbalist.....	5
26	Do you use homemade oral rehydration solution?	Yes.....	1
		No.....	2
27	If yes, how do you prepare it?	1 tea spoonful of salt, 8 tea spoonful of sugar in one litre of water.....	1
		2 tea spoonful of salt, 4 tea spoonful of sugar in one litre of water.....	2
		8 teaspoonful of salt to 8 tea spoonful of sugar.....	3
		Don't know.....	4
28	Would you give ORS at every watery stool?	Yes.....	1
		No.....	2
		Don't know.....	3
29	Do you consider diarrhoea to be severe when the stool is bloody?	Yes.....	1
		No.....	2
		Don't know.....	3
30	Do you consider diarrhoea to be severe when the child has sunken fontanel?	Yes.....	1
		No.....	2
		Don't know.....	3
31	Do you consider diarrhoea to be severe when child cries but with no tears?	Yes.....	1
		No.....	2

No	Questions and filters	Coding categories	Skip to
32	What action would you take if your child's diarrhoea gets worse?	Try home remedies.....1 Take child to traditional healers.....2 Take child for medical care at a clinic.....3 Other, specify.....4	

APPENDIX TWO: INFORMED CONSENT AND PARTICIPANTS DECLARATION

Informed Consent

Date:
Good day mothers

My name is: Adeleke Adekunle Isaac, and I am a student currently enrolled for a Master's degree in Public Health, Howard College Campus, at the University of KwaZulu-Natal (UKZN), Durban in South Africa. The reason I came here is to ask some questions related to child Health, in order to understand your level of knowledge, your attitudes and practices in the management of diarrhoea in under five year old children. This research process forms part of my Master's thesis entitled:

"Maternal knowledge, attitudes and practices towards prevention and management of child diarrhoea in urban and rural Maseru, Lesotho, 2016"

The study aimed to understand better the knowledge, attitudes and practices of mothers in the management of diarrhoea in relation to where they live. This will help in understanding the different ways that mothers prevent and manage diarrhoea in children less than five years old in the urban and rural settings.

The research will require about 500 participants, with a questionnaire containing about 36 questions required to be completed by the participants with the assistance of researchers. Participants will be required to answer the questionnaire provided to them after they have fully agreed to do so voluntarily.

The result of the research will help the participants and the government of Lesotho and developing countries at large, in knowing the approach to apply in providing basic information and effective health education components, to strengthen health education programmes for healthcare facilities in the management of diarrhoea. Each participant will be allocated a number, therefore the names will not be revealed.

There is no material or financial benefits attached to participating in this research study, and your participation is entirely voluntary. Please note that any participant can withdraw from this study at any time, there will not be any loss of services she is entitled to.

The information obtained from the questionnaire will be treated in a confidential manner, and will be safely stored in a locker at the School of Public Health, University of KwaZulu-Natal.

Should you need further clarity or have any questions regarding this research study, please contact me or my research supervisor.

Researcher:

Adeleke Adekunle

Research Supervisor

Dr Tsholofelo Mhlaba

Your participation is much appreciated, thank you.
BREC ETHICAL APPROVAL NUMBER (BE588/16)
BIOMEDICAL RESEARCH ETHICS ADMINISTRATION
Research Office, Westville Campus
Govan Mbeki Building
University of KwaZulu-Natal
Private Bag X 54001, Durban, 4000
KwaZulu-Natal, SOUTH AFRICA
Tel: 27 31 2602486 - Fax: 27 31 2604609

DECLARATION

I..... hereby declare that I am fully aware of the contents of this Informed Consent Form and the nature of this research project. I fully agree to participate in this research project as a volunteer, and, therefore, I have the right to refuse to answer any questions as per my discretion.

I also have the right to withdraw from this research study at any point, should I wish to do so, and my actions will not disadvantage me in any way. I will not receive any payment for participating in the research.

Signature of Participant.....

Witness.....

Date.....

LESOTHO MINISTRY OF HEALTH APPROVAL



LESOTHO

Ministry of Health
PO Box 514
Maseru 100

REF: ID113-2016

Date: 01 November 2016

To
Adeleke Adekunle Isaac
Student number 2150 73608
Masters of PH candidate
University of KWAZULU-NATAL, RSA

Category of Review:

- Initial Review
- Continuing Annual Review
- Amendment/Modification
- Reactivation
- Serious Adverse Event
- Other _____

Dear Adeleke,

RE: Maternal knowledge, attitudes and practices towards prevention and management in child diarrhea in urban and rural Maseru, Lesotho, 2016

This is to inform you that on 20 October 2016 the Ministry of Health Research and Ethics Committee reviewed and **APPROVED** the modifications of above named protocol and hereby authorizes you to continue the study according to the activities and population specified in the protocol. Departure from the approved protocol will constitute a breach of this permission.

This approval includes review of the following attachments:

- Protocol dated 08 September 2016
 - English & Sesotho consent forms
 - Data collection forms in Sesotho
 - Data collection forms in English
 - Participant materials *[insert types, versions, dates]*
 - Other materials: The letter of recommendation from KWAZULU-NATAL University
- This approval is **VALID** until 24 October 2017.

Please note that an annual report and request for renewal, if applicable, must be submitted at least 6 weeks before the expiry date.

All serious adverse events associated with this study must be reported promptly to the MOH Research and Ethics Committee. Any modifications to the approved protocol or consent forms must be submitted to the committee prior to implementation of any changes.

We look forward to receiving your progress reports and a final report at the end of the study. If you have any questions, please contact the Research and Ethics Committee at rcumoh@gmail.com (or) 22226317.

Sincerely,

Dr. Nyane Letsie
Director General Health Services

Mrs. V. T. Lenana
Co-chairperson NH-IRB



*and Domesticity
and Tlali
Please assist*

BREC (BIOMEDICAL RESEARCH ETHICS COMMITTEE) APPROVAL



26 January 2017

Dr Al Adeleke (215073608)
Discipline of Public Health
School of Nursing and Public Health Medicine
Health Sciences
docleke@gmail.com

Protocol: Maternal knowledge, attitudes and practices towards prevention and management of child diarrhoea in urban and rural Maseru, Lesotho, 2016.

Degree: MPH

BREC reference number: BE588/16

EXPEDITED APPLICATION

A sub-committee of the Biomedical Research Ethics Committee has considered and noted your application received on 21 October 2016.

The study was provisionally approved pending appropriate responses to queries raised. Your response received on 18 January 2017 to BREC letter dated 14 December 2016 have been noted by a sub-committee of the Biomedical Research Ethics Committee. The conditions have now been met and the study is given full ethics approval and may begin as from 26 January 2017.

This approval is valid for one year from 26 January 2017. To ensure uninterrupted approval of this study beyond the approval expiry date, an application for recertification must be submitted to BREC on the appropriate BREC form 2-3 months before the expiry date.

Any amendments to this study, unless urgently required to ensure safety of participants, must be approved by BREC prior to implementation.


Your acceptance of this approval denotes your compliance with South African National Research Ethics Guidelines (2015), South African National Good Clinical Practice Guidelines (2006) (if applicable) and with UKZN BREC ethics requirements as contained in the UKZN BREC Terms of Reference and Standard Operating Procedures, all available at <http://research.ukzn.ac.za/Research-Ethics/Biomedical-Research-Ethics.aspx>.

BREC is registered with the South African National Health Research Ethics Council (REC-290408-009). BREC has US Office for Human Research Protections (OHRP) Federal-wide Assurance (FWA 678).

The sub-committee's decision will be RATIFIED by a full Committee at its next meeting taking place on 14 February 2017.

We wish you well with this study. We would appreciate receiving copies of all publications arising out of this study.

Yours sincerely


Professor Joyce Tsoka-Gwegweni
Chair: Biomedical Research Ethics Committee

cc supervisor: mhlaba@ukzn.ac.za
cc postgraduate administrator: anumugamc@ukzn.ac.za

Biomedical Research Ethics Committee
Professor J Tsoka-Gwegweni (Chair)
Westville Campus, Govan Mbeki Building
Postal Address: Private Bag X54001, Durban 4000

Telephone: +27 (0) 31 260 2485 Facsimile: +27 (0) 31 260 4809 Email: brec@ukzn.ac.za

© 2019 Adeleke and Mhlaba; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/49177>